

REPORT DOCUMENTATION

AD-A281 282

-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing the collection of information, gathering and maintaining the data needed, and completing and reviewing the collection of information, including suggestions for reducing this burden, to Washington, Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Project, Washington, DC 20503.

isting data sources
other aspect of this
rts, 1215 Jefferson
0503.

| | | |
|--|--------------------------------|---|
| 1. AGENCY USE ONLY (Leave blank) | 2. REPORT DATE May 13, 1994 | 3. REPORT TYPE AND DATES COVERED Final 1 Nov 92-15 Apr 94 |
| 4. TITLE AND SUBTITLE Conformation of Membrane Proteins: Bacteriorhodopsin | | 5. FUNDING NUMBERS DAAH04-93-G-0007 |
| 6. AUTHOR(S) Gerald D. Fasman | | 8. PERFORMING ORGANIZATION REPORT NUMBER |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Brandeis University Waltham, Massachusetts 02254-9110 | | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211 | | 10. SPONSORING/MONITORING AGENCY REPORT NUMBER ARO 29714.1-LS |
| 11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation. | | |
| 12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited. | | 12b. DISTRIBUTION CODE |

13. ABSTRACT (Maximum 200 words)

Bacteriorhodopsin, from the purple membrane (PM) of *Halobacterium halobium*, was chemically modified with methoxypolyethylene glycol (MW = 5000) succinimidyl carbonate. The polyethylene glycol-bacteriorhodopsin (m-PEG-SC-BR33) conjugate, containing one PEG chain, was water soluble. The secondary structure of the conjugate in water appeared partially denatured but was shown to contain α -helical segments

(continued on reverse side)

| | | | |
|---|---|--|--------------------------------------|
| 14. SUBJECT TERMS Membrane Proteins Bacteriorhodopsin | | | 15. NUMBER OF PAGES |
| | | | 16. PRICE CODE |
| 17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED | 18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED | 19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED | 20. LIMITATION OF ABSTRACT UL |

NSN 7540-01-280-5500

Standard Form 298 (Rev 2-89)
Prescribed by ANSI Std Z39-18
298-102

94 7 11 0 2 6

94-21110

by circular dichroism (CD) spectroscopy. The isolated bacteriorhodopsin conjugate, with added retinal, was refolded in a mixed detergent-lipid micelle and had an absorption maximum at 555 nm. The refolded conjugate was transferred into vesicles which pumped protons, upon illumination, as efficiently as did native BR. Modification of the PM with methoxy-polyethylene glycol did not alter the native structure or inhibit proton pumping, and therefore it is suggested that the glycol polymer is present as a covalently linked moiety to residues unnecessary for proton pumping and proper folding. The site of attachment of mPEG was determined to be either at Lys 129 or Lys 159, with position Lys 129 the most probable site of attachment. The m-PEG-SC-BR33 could be stepwise refolded to the native conformation by the addition of trifluoroethanol to lower the dielectric constant, simulating the insertion of the BR into the phospholipid bilayer.

CONFORMATION OF MEMBRANE PROTEINS: BACTERIORHODOPSIN

FINAL REPORT

GERALD D. FASMAN

MAY 13, 1994

U.S. ARMY RESEARCH OFFICE
CHEMICAL AND BIOLOGICAL SCIENCES DIVISION

GRANT NUMBER DAAH04-93-G-0007

APPROVED FOR PUBLIC RELEASE

DISTRIBUTION UNLIMITED

| | |
|---------------------|---|
| Accession For | |
| NTIS | CRA&I <input checked="" type="checkbox"/> |
| DTIC | TAB <input checked="" type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification _____ | |
| By _____ | |
| Distribution / | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A-1 | |

THE VIEWS, OPINIONS, AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHOR(S) AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION, POLICY, OR DECISION, UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.

4. REPORT

A. STATEMENT OF PROBLEM

Bacteriorhodopsin (BR) is an integral membrane protein (248 amino acids) which catalyzes the light-induced proton translocation across the membrane of *Halobacterium halobium*. Research was directed toward: (1) the elucidation of the folding path of BR as it attains its final conformation in the membrane, and, (2) to facilitate goal #1, a water-soluble derivative, a methoxy polyethylene glycol derivative (m-PEG-BR) will be synthesized.

B. SUMMARY OF RESULTS

Bacteriorhodopsin, from the purple membrane (PM) of *Halobacterium halobium*, was chemically modified with methoxypolyethylene glycol (MW = 5000) succinimidyl carbonate. The polyethylene glycol-bacteriorhodopsin (m-PEG-SC-BR33) conjugate, containing one PEG chain, was water soluble. The secondary structure of the conjugate in water appeared partially denatured but was shown to contain α -helical segments by circular dichroism (CD) spectroscopy. The isolated bacteriorhodopsin conjugate, with added retinal, was refolded in a mixed detergent-lipid micelle and had an absorption maximum at 555 nm. The refolded conjugate was transferred into vesicles which pumped protons, upon illumination, as efficiently as did native BR. Modification of the PM with methoxy-

polyethylene glycol did not alter the native structure or inhibit proton pumping, and therefore it is suggested that the glycol polymer is present as a covalently linked moiety to residues unnecessary for proton pumping and proper folding. The site of attachment of mPEG was determined to be either at Lys 129 or Lys 159, with position Lys 129 the most probable site of attachment. The m-PEG-SC-BR33 could be stepwise refolded to the native conformation by the addition of trifluoroethanol to lower the dielectric constant, simulating the insertion of the BR into the phospholipid bilayer.

C. PUBLICATIONS

1. "The Refolding and Proton Pumping Activity of a Polyethylene Glycol-Bacteriorhodopsin Water-Soluble Conjugate," G. Fasman and G. Sirokmán. Sixth Symp. of the Protein Society, 1992.
2. "Refolding and Proton Pumping Activity of a Polyethylene Glycol-Bacteriorhodopsin Water-Soluble Conjugate," G. Sirokmán and G.D. Fasman. *Protein Science* 2, 1161-1170 (1993).
3. "Crystallizing Membrane Proteins. A New Successful Method," G.D. Fasman and G. Sirokmán, *FASEB J.* 8, Abst. 1113 (1994), Amer. Soc. Biochem. and Mol. Biol., 1994 Meeting, Washington, D.C.

D. PARTICIPATING SCIENTIFIC PERSONNEL

Gerald D. Fasman

Géza Sirokmán

5. NO INVENTIONS